

## Cholestin<sup>®</sup>

Elson CE, Qureshi AA. Coupling the cholesterol- and tumor-suppressive actions of palm oil to the impact of its minor constituents on 3-hydroxy-3-methylglutaryl coenzyme A reductase activity. *Prostaglandins Leukot Essent Fatty Acids* 1995; 52(2-3):205-207.

Jenkins DJA, Kendall CWC, Marchie A, Faulkner D, Vidgen E, Lapsley KG et al. The effect of combining plant sterols, soy protein, viscous fibers, and almonds in treating hypercholesterolemia. *Metabolism* 2003; 52(11):1478-1483.

Kurowska EM, Manthey JA, Casaschi A, Theriault AG. Modulation of HepG2 cell net apolipoprotein B secretion by the citrus polymethoxyflavone, tangeretin. *Lipids* 2004; 39(2):143-151.

Kurowska EM, Manthey JA. Hypolipidemic effects and absorption of citrus polymethoxylated flavones in hamsters with diet-induced hypercholesterolemia. *J Agric Food Chem* 2004; 52(10):2879-2886.

Kurowska EM, Manthey JA. Regulation of lipoprotein metabolism in HepG2 cells by citrus flavonoids. *Adv Exp Med Biol* 2002; 505:173-179.

Pearce BC, Parker RA, Deason ME, Dischino DD, Gillespie E, Qureshi AA et al. Inhibitors of cholesterol biosynthesis. 2. Hypocholesterolemic and antioxidant activities of benzopyran and tetrahydronaphthalene analogues of the tocotrienols. *J Med Chem* 1994; 37(4):526-541.

Pearce BC, Parker RA, Deason ME, Qureshi AA, Wright JJ. Hypocholesterolemic activity of synthetic and natural tocotrienols. *J Med Chem* 1992; 35(20):3595-3606.

Peffley DM, Gayen AK. Plant-derived monoterpenes suppress hamster kidney cell 3-hydroxy-3-methylglutaryl coenzyme a reductase synthesis at the post-transcriptional level. *J Nutr* 2003; 133(1):38-44.

Peterson DM, Qureshi AA. Effects of tocols and  $\beta$ -glucan on serum lipid parameters in chickens. *J Sci Food Agric* 1997; 73(4):417-424.

Qureshi AA, Bradlow BA, Brace L, Manganello J, Peterson DM, Pearce BC et al. Response of hypercholesterolemic subjects to administration of tocotrienols. *Lipids* 1995; 30(12):1171-1177.

Qureshi AA, Bradlow BA, Salser WA, Brace LD. Novel tocotrienols of rice bran modulate cardiovascular disease risk parameters of hypercholesterolemic humans. *J Nutr Biochem* 1997; 8(5):290-298.

Qureshi AA, Burger WC, Peterson DM, Elson CE. The structure of an inhibitor of

cholesterol biosynthesis isolated from barley. *J Biol Chem* 1986; 261(23):10544-10550.

Qureshi AA, Lehmann JW, Peterson DM. Amaranth and its oil inhibit cholesterol biosynthesis in 6-week-old female chickens. *J Nutr* 1996; 126(8):1972-1978.

Qureshi AA, Mo HB, Packer L, Peterson DM. Isolation and identification of novel tocotrienols from rice bran with hypocholesterolemic, antioxidant, and antitumor properties. *J Agric Food Chem* 2000; 48(8):3130-3140.

Qureshi AA, Peterson DM. The combined effects of novel tocotrienols and lovastatin on lipid metabolism in chickens. *Atherosclerosis* 2001; 156(1):39-47.

Qureshi AA, Qureshi N, Hasler-Rapacz JO, Weber FE, Chaudhary V, Crenshaw TD et al. Dietary tocotrienols reduce concentrations of plasma cholesterol, apolipoprotein B, thromboxane B<sub>2</sub>, and platelet factor 4 in pigs with inherited hyperlipidemias. *Am J Clin Nutr* 1991; 53(4 Suppl):1042S-1046S.

Qureshi AA, Qureshi N, Wright JJ, Shen Z, Kramer G, Gapor A et al. Lowering of serum cholesterol in hypercholesterolemic humans by tocotrienols (palmvitee). *Am J Clin Nutr* 1991; 53(4 Suppl):1021S-1026S.

Qureshi AA, Salser WA, Parmar R, Emeson EE. Novel tocotrienols of rice bran inhibit atherosclerotic lesions in C57BL/6 apoE-deficient mice. *Journal of Nutrition* 2001; 131(10):2606-2618.

Qureshi AA, Sami SA, Salser WA, Khan FA. Dose-dependent suppression of serum cholesterol by tocotrienol-rich fraction (TRF25) of rice bran in hypercholesterolemic humans. *Atherosclerosis* 2002; 161(1):199-207.

Qureshi N, Qureshi AA. Tocotrienols, Novel Hypocholesterolemic Agents with Antioxidant Properties. In: Packer L, Fuchs J, editors. *Vitamin E in Health and Disease*. New York: Marcel Dekker, 1993: 247-267.

Roche HM, Gibney MJ. Postprandial triacylglycerolaemia: the effect of low-fat dietary treatment with and without fish oil supplementation. *Eur J Clin Nutr* 1996; 50(9):617-624.

Simopoulos AP. Summary of the NATO advanced research workshop on dietary omega 3 and omega 6 fatty acids: biological effects and nutritional essentiality. *J Nutr* 1989; 119(4):521-528.

Theriault A, Chao JT, Wang Q, Gapor A, Adeli K. Tocotrienol: a review of its therapeutic potential. *Clin Biochem* 1999; 32(5):309-319.

Theriault A, Wang Q, Gapor A, Adeli K. Effects of gamma-tocotrienol on ApoB synthesis, degradation, and secretion in HepG2 cells. *Arterioscler Thromb Vasc Biol*

1999; 19(3):704-712.

Wang Q, Theriault A, Gapor A, Adeli K. Effects of tocotrienol on the intracellular translocation and degradation of apolipoprotein B: possible involvement of a proteasome independent pathway. *Biochem Biophys Res Commun* 1998; 246(3):640-643.

Whitman SC, Kurowska EM, Manthey JA, Daugherty A. Nobiletin, a citrus flavonoid isolated from tangerines, selectively inhibits class A scavenger receptor-mediated metabolism of acetylated LDL by mouse macrophages. *Atherosclerosis* 2005; 178(1):25-32.